

Two new yellow-legged species of *Hybos* Meigen from Hainan, China (Diptera, Hybotidae)

Ding YANG

Department of Entomology, China Agricultural University, Beijing 100094, China.

E-mail: dyangcau@yahoo.com.cn

Two new yellow-legged species of *Hybos* Meigen from Hainan, China (Diptera, Hybotidae). - Six yellow-legged species of the genus *Hybos* are reported from Hainan. The following two new species are described, illustrated and compared with morphologically similar species: *Hybos bawanglingensis* sp. n. and *H. hainanensis* sp. n. A key to the six species of the genus from Hainan is presented for the first time.

Keywords: Diptera - Hybotidae - *Hybos* - new species - Hainan.

INTRODUCTION

Within the Hybotidae *Hybos* Meigen is characterized by the following traits: long spine-like proboscis, anal cell longer than basal cell, Rs rather short, R_{4+5} and M_1 divergent apically (Chvála, 1983; Yang & Yang, 2004). It is distributed worldwide with 164 known species (Yang *et al.*, 2007). The species of *Hybos* from China were revised by Yang & Yang (2004) who mentioned 85 known species. The major references dealing with *Hybos* in the Old World are Brunetti (1920), Frey (1953), Smith (1965), Saigusa (1963, 1965), Chvála (1983), Yang & Yang (2004), and Yang *et al.* (2006).

Hainan Island with a tropical climate belongs to South China. It is famous for its rich biodiversity. Up to now, only two *Hybos* species, *H. jianfengensis* Yang, Yang & Hu, 2002 and *H. particularis* Yang, Yang & Hu, 2002, have been described from this region (Yang *et al.*, 2002; Yang & Yang, 2004). In the present paper, four yellow-legged species of *Hybos* are added to the fauna of Hainan. Two species are described as new to science. A key to the species of the genus from Hainan is presented for the first time.

MATERIAL AND METHODS

The specimens for this study are deposited in the following collections:

CAU Entomological Museum of China Agricultural University, Beijing, China.

MHNG Muséum d'histoire naturelle, Genève, Switzerland.

Basic terminology follows McAlpine (1981) and Steyskal & Knutson (1981). The following abbreviations are used: acr = acrostichal, ad = anterodorsal, av = anteroventral, d = dorsal, dc = dorsocentral, h = humeral, npl = notopleural, oc = ocellar, pd = posterodorsal, presc = prescutellar, psa = postalar, pv = posteroventral, v = ventral.

KEY TO SPECIES (MALES) OF *HYBOS* FROM HAINAN

1. Legs mostly yellow; arista bare or pubescent 2
 Legs mostly or entirely black; arista bare 5
2. Thoracic pleuron black; scutellum black 3
 Thoracic pleuron yellow; scutellum yellow
 *flaviscutellum* Yang & Yang, 1986
3. Hind tibia with 1 strong ad at middle; arista pubescent at least on basal 3/4 . 4
 Hind tibia without distinct ad; arista bare *hainanensis* sp. nov.
4. Left and right surstyli hook-like; hypandrium with a large apical incision *serratus* Yang & Yang, 1992
 Left and right surstyli not hook-like; hypandrium with a narrow cleft apically *bawanglingensis* sp. nov.
5. Legs entirely black; wing pale grayish brown
 *jianfengensis* Yang, Yang & Hu, 2002
 Legs partly yellow; wing hyaline *particularis* Yang, Yang & Hu, 2002

TAXONOMIC PART

Hybos bawanglingensis sp. n.

Figs 1-4

MATERIAL: Holotype male, Hainan, Bawangling, Donger, 25. V. 2007, Junhua Zhang (CAU).


ETYMOLOGY: The species is named after the type locality Bawangling.

DIAGNOSIS: Legs yellow except tarsomeres 3-5 black. Hind tibia with 1 ad at middle. Hypandrium with a narrow cleft apically.

DESCRIPTION: MALE. Body length 4.2 mm, wing length 3.8 mm.

Head: black, pale gray microtrichose. Eyes dark brown but enlarged upper facets brownish yellow, contiguous on frons. Setae and setulae on head black; ocellar tubercle weak, with one pair of long oc and 2 very short posterior setulae. Antenna black; scape bare; pedicel with a circle of subapical setulae; first flagellomere black without dorsal setulae; arista black, short pubescent except apical 1/4 thin and bare. Proboscis black; palpus black with 1 long ventral setula apically.

Thorax: black, pale grey microtrichose. Setae and setulae on thorax black; postpronotal lobe with black setulae; h absent, 2 npl (anterior npl short), acr in about 6 more or less regular rows, 1 distinct posterior dc and 8 setulae anteriorad, 1 long presc, 1 long psa; scutellum with 6 marginal setulae and 2 long subapical setae. Legs yellow; hind femur dark yellow dorsally, hind knee dark brown, hind tibia dark yellow basally; tarsomeres 3-5 dark brown. Setae and setulae on legs black. Fore femur 1.1 times as wide as mid femur, with one row of setulae-like pv nearly as long as femur thickness; hind femur 1.9 times as wide as mid femur, with 3 preapical anterior setae, and with one row of 9 v and 6 av which are spine-like and inserted on weak tubercles. Fore tibia with 1 long d at middle, apically with 1 long ad; mid tibia with 2 long ad on basal half and 1 very long av at middle, apically with 5 setae of which 1 av is very long; hind tibia with 1 long ad at middle, apically with 1 av and pd. Fore tarsomere 1 with 1 v at base; mid tarsomere 1 with 2 v on basal 2/5, apically with 1 ad and 1 pv; hind tarso-



FIGS 1-4

Hybos bawanglingensis sp. n. (male). (1) genitalia, dorsal view. (2) right surstylus. (3) left surstylus. (4) hypandrium, ventral view.

mere 1 with 2 v at base. Wing slightly tinged with grayish; stigma dark brown; veins dark brown, R_{4+5} and M_1 divergent apically. Squama dark yellow, bordered with pale setulae. Halter brownish.

Abdomen: nearly straight, black, grey microtrichose. Setulae on abdomen black. Male genitalia (Figs 1-4): left epandrial lobe narrow, left surstylus with a short and a long process closely located; right epandrial lobe wide, right surstylus with a long spine-like process strongly curved inward; hypandrium distinctly longer than wide, basally wide and nearly truncate, apically narrow and with a narrow cleft.

FEMALE: Unknown.

DISTRIBUTION: China (Hainan).

REMARKS: The new species is very similar to *Hybos gutianshanus* Yang & Yang from Zhejiang and Guangdong, but can be easily separated from the latter by having the hind femur with 3 preapical anterior setae, the hind tibia with 1 ad at middle, and the hypandrium narrowly cleft apically with the right lobe long and narrow apically. In *H. gutianshanus*, the hind femur has 4 preapical anterior setae, the hind tibia has 2 ad at middle, and the hypandrium is widely cleft apically with the right lobe wide and obtuse apically (Yang & Yang, 2004).

Hybos hainanensis sp. n.

Figs 5-8

MATERIAL: Holotype male, Hainan, Wuzhishan, Jingguantai-Zhumanguzhai, 18. V. 2007, Kuiyan Zhang (CAU). Paratypes: 1 male, 1 female, Hainan, Wuzhishan, Shuimanxiang, 15. V. 2007, Jie Zeng (CAU). 6 males, 6 females, Hainan, Diaoluoshan, Baishuiling, 1-2. VI. 2007, Jingxian Liu (CAU, MHNG).

ETYMOLOGY: The species is named after the type locality Hainan.

DIAGNOSIS: Arista bare. Legs yellow except hind knee and base of hind tibia blackish and tarsomeres 3-5 dark brown. Left epandrial lobe with 4 long strong spines

DESCRIPTION: MALE. Body length 3.0-3.8 mm, wing length 2.8-3.8 mm.

Head: black, pale gray microtrichose. Eyes dark brown but enlarged upper facets brownish yellow, contiguous on frons. Setae and setulae on head black; ocellar tubercle weak, with one pair of long oc and 2 very short posterior setulae. Antenna black; scape bare; pedicel with a circle of subapical setulae; first flagellomere black without long seta (but with 1 subapical long seta in 1 paratype); arista black, bare. Proboscis blackish; palpus black with 1 long ventral setula apically.


Thorax: black, pale gray microtrichose. Setae and setulae on thorax black; postpronotal lobe with black setulae; h absent, 2 npl (anterior npl short), acr in about 6 more or less regular rows, 1 distinct posterior dc and 8 setulae anteriorly, 1 long presc, 1 long psa; scutellum with 6 marginal setulae and 2 long subapical setae. Legs yellow, but base of hind tibia including knee blackish and tarsomeres 3-5 dark brown. Setae and setulae on legs black. Fore femur 1.2 times as wide as mid femur, with one row of long setula-like pv distinctly longer than femur thickness; mid femur with one row of setula-like av and pv nearly as long as femur thickness; hind femur 2.7 times as wide as mid femur, with 3 preapical dorsal setae, and with one row of 15 v and 7 av which are spine-like and inserted on weak tubercles. Fore tibia with 1 long ad at middle and one row of very long setula-like pv (longer than pv on fore femur), apically with 1 long ad; mid tibia with 1 very long ad at basal 1/3, 1 short ad at apical 1/3 and 1 very long av at middle, apically with 4 setae (1 av very long); hind tibia without ad but with a long setula at middle, apically with 1 av and 1 pd. Fore tarsomere 1 with long setula-like pv; mid tarsomere 1 with 1 v at basal 1/3; hind tarsomeres 1 with 2 very short ventral spines on basal 1/3. Wing hyaline; stigma dark brown; veins dark brown, R_{4+5} and M_1 divergent apically. Squama dark yellow, bordered with pale setulae. Halter pale yellow.

Abdomen: nearly straight, black, grey microtrichose. Setulae on abdomen black. Male genitalia (Figs 5-8): left epandrial lobe wide with one row of 4 long strong spines, left surstylus curved, more or less tapering toward tip, with 1 seta at extreme tip; right epandrial lobe wide, right surstylus deeply bifurcated apically; hypandrium much longer than wide, basally slightly narrow, apically furcated.

FEMALE: Body length 3.0-3.9 mm, wing length 3.0-3.7 mm. Similar to male.

DISTRIBUTION: China (Hainan).

REMARKS: The new species is very particular. It can be easily separated from other known species of the genus by the left epandrial lobe with 4 long strong spines.



FIGS 5-8

Hybos hainanensis sp. n. (male). (5) genitalia, dorsal view. (6) right surstylus. (7) left surstylus. (8) hypandrium, ventral view.

Hybos flaviscutellum Yang & Yang, 1986.

Hybos flaviscutellum Yang & Yang, 1986: 81. Type locality: China: Guangxi, Longsheng.

MATERIAL: 1 male, Hainan, Bawangling, Donger, 24. V. 2007, Junhua Zhang (CAU).

DISTRIBUTION: China (Zhejiang, Guangxi, Hainan).

Hybos serratus Yang & Yang, 1992.

Hybos serratus Yang & Yang, 1992: 1089. Type locality: China: Sichuan, Xichang.

MATERIAL: 1 male, Hainan, Bawangling, Donger, 24. V. 2007, Kuiyan Zhang (CAU).

DISTRIBUTION: China (Henan, Sichuan, Guizhou, Guangxi, Hainan).

ACKNOWLEDGEMENTS

My sincere thanks are due to Dr. Junhua Zhang and Ms. Kuiyan Zhang (China Agricultural University, Beijing) and Mr. Jingxian Liu and Ms. Jie Zeng (South Agricultural University, Guangzhou) for collecting the specimens, and to Dr. B. Merz (Muséum d'histoire naturelle, Genève) for reviewing an earlier draft of this manuscript. The research was funded by the National Natural Science Foundation of China (No. 30770259, No. 30225009) and the Ministry of Science and Technology of the People's Republic of China (MOST grant no. 2006FY110500).

REFERENCES

- BRUNETTI, E. 1920. Diptera Brachycera Vol. 1. *The Fauna of British India, including Ceylon & Burma*. London, 401 pp.
- CHVÁLA, M. 1983. The Empidoidea (Diptera) of Fennoscandia and Denmark. II. General Part. The families Hybotidae, Atelestidae and Microphoridae. *Fauna Entomologica Scandinavica* 12: 279 pp. Scandinavian Science Press, Copenhagen.
- FREY, R. 1953. Studien über ostasiatische Dipteren. II. Hybotinae, Ocydromiinae, *Hormopeza* Zett. *Notulae Entomologicae* 33: 57-71.
- MCALPINE, J. F. 1981. Morphology and terminology - adults. (pp. 9-63). In: MCALPINE, J. F., PETERSON, B. V., SHEWELL, G. E., TESKEY, H. J., VOCKEROTH, J. R. & WOOD, D. M. (eds). *Manual of Nearctic Diptera*. Vol. 1. *Research Branch, Agriculture Canada, Monograph* 27: 1-674.
- SAIGUSA, T. 1963. Systematic studies of the genus *Hybos* in Japan I. New species with yellowish legs (Diptera, Empididae). *Sieboldia* 3(1): 97-104.
- SAIGUSA, T. 1965. Studies on the Formosan Empididae collected by Professor T. Shirôzu (Diptera, Brachycera). *Special Bulletin of Lepidoptera Society in Japan* 1: 180-196.
- SMITH, K. G. V. 1965. Diptera from Nepal: Empididae. *Bulletin of the British Museum (Natural History), Entomology* 17(2): 61-112.
- STEYSKAL, G. C. & KNUTSON, L. V. 1981. 47. Empididae (pp. 607-624). In: MCALPINE, J. F., PETERSON, B. V., SHEWELL, G. E., TESKEY, H. J., VOCKEROTH, J. R. & WOOD, D. M. (eds). *Manual of Nearctic Diptera*. Vol. 1. *Research Branch, Agriculture Canada, Monograph* 27: 1-674.
- YANG, C. & YANG, D. 1986. Fourteen new species of dance flies from Fujian and Hainan (Diptera: Empididae). *Wuyi Science Journal* 6: 75-88.
- YANG, D., MERZ, B. & GROOTAERT, P. 2006. New yellow-legged *Hybos* from Nanling, Guangdong, China (Diptera, Empidoidea, Hybotidae). *Revue suisse de Zoologie* 113(4): 797-806.
- YANG, D. & YANG, C. 1992. Diptera: Empididae (pp. 1089-1097). In: CHEN, S. (ed.). *Insects of the Hengduan Mountains Region*, Vol. 2: 867-1547. Science Press, Beijing.
- YANG, D. & YANG, C. 2004. *Fauna Sinica Insecta* Vol. 34. Diptera, Empididae, Hemerodromiinae Hybotinae. Science Press, Beijing, 329 pp.
- YANG, D., YANG, C. & HU, X. 2002. Diptera: Empididae (pp. 733-740). In: HUANG, F. (ed.). *Forestry Insects of Hainan*. Science Press, Beijing, 1064 pp.
- YANG, D., ZHANG, K. Y., YAO, G. & ZHANG, J. H. 2007. *World catalog of Empididae (Insecta: Diptera)*. China Agricultural University Press, Beijing, 599 pp.